

Name: \_\_\_\_\_

Discussion Section: \_\_\_\_\_

---

**Solutions should show all of your work, not just a single final answer.**

---

## 4.7: Optimization Problems

---

1. A closed box (top, bottom, and all four sides) needs to be constructed to have a volume of  $9\text{ m}^3$  and a base whose width is twice its length. See Figure 1.




Figure 1: A box

Use calculus to determine the dimensions (length, width, height) of such a box that uses the least amount of material. Justify why your answer corresponds to a minimum, not a maximum.

2. We want to find the points on  $y = x^2$  that are closest to  $(0, 3)$ .




Figure 2: Distance to  $(0, 3)$  on  $y = x^2$ .

- (a) For each point  $(x, x^2)$  on the parabola, find a formula for its distance to  $(0, 3)$ . Call this distance  $D(x)$ . (See Figure 5.)
- (b) Let  $f(x) = D(x)^2$ , which is the *squared distance* between  $(x, x^2)$  and  $(0, 3)$ . Finding where  $D(x)$  is minimal is the same as finding where  $f(x)$  is minimal. Determine all  $x$  where  $f(x)$  has an absolute minimum. The points  $(x, x^2)$  for such  $x$  are the closest points to  $(0, 3)$  on  $y = x^2$ .

3. Three line segments of length 1 are joined together at endpoints to form a base and the legs of an isosceles trapezoid, as in Figure 3. Let  $\theta$  in  $(0, \pi/2)$  be the common angle measurement between the legs and the line passing through the base of length 1. We want to find the angle  $\theta$  that maximizes the area of the trapezoid.




Figure 3: An isosceles trapezoid with base and legs of length 1.

- (a) Compute the area  $A(\theta)$  of the trapezoid. The general area formula for a trapezoid is  $\frac{1}{2}h(b_1 + b_2)$ , where  $h$  is the height and  $b_1$  and  $b_2$  are the lengths of the bases. (Hint: Break up the trapezoid into a rectangle with two right triangles at both ends. Use trigonometry to compute the height and the length of the longer base in terms of  $\theta$ .)
- (b) Find all solutions to  $A'(\theta) = 0$  with  $0 < \theta < \pi/2$ . (The answer is *not*  $\pi/4 = 45^\circ$ .)
- (c) Verify that the area  $A(\theta)$  is a maximum, not a minimum, at the angle found in part (b) and compute this maximum area.

---

## 4.8: Newton's Method

---

4. Apply Newton's method to estimate the solution of  $x^3 - x - 1 = 0$  by taking  $x_1 = 1$  and finding the least  $n$  such that  $x_n$  and  $x_{n+1}$  agree to three digits after the decimal point.

5. The number  $\pi$  is a solution of  $\sin x = 0$  close to 3 (see Figure 4). You will use Newton's method for  $\sin x = 0$  to create numerical estimates for  $\pi$ .




Figure 4: Graph of  $y = \sin x$ .

- (a) Write out the recursion for Newton's method used to solve  $\sin x = 0$ .

(b) Using Newton's method for  $\sin x = 0$  with  $x_1 = 3$ , find the first  $n$  for which  $x_n$  and  $x_{n+1}$  agree to 5 digits after the decimal point. (Use radians, *not* degrees!)

(c) For the  $n$  you found in part (b), to how many digits after the decimal point does  $x_n$  actually agree with  $\pi$ ?

6. In Figure 5 is the graph of  $f(x) = \ln(x) - 1$  for  $0 < x < 4$ . It crosses the  $x$ -axis at  $x = e$ . You will use Newton's method for  $f(x) = 0$  to create numerical estimates for  $e$ .




Figure 5: Graph of  $y = \ln(x) - 1$ .

- (a) Using Newton's method for the equation  $\ln(x) - 1 = 0$  with  $x_1 = 1$ , tabulate  $x_n$  to find the first  $n$  for which  $x_n$  and  $x_{n+1}$  agree to 5 digits after the decimal point.
- (b) For the  $n$  you found in part (a), to how many digits after the decimal point does  $x_n$  actually agree with  $e$ ?

Answers to Selected Problems:

1. The dimension are length  $= x = \frac{3}{2}$ , width  $= 2x = 3$ , and height  $= \frac{9}{2(3/2)^2} = 2$ .
2. (a)  $D(x) = \sqrt{x^2 + (x^2 - 3)^2}$   
(b)  $f(x)$  has absolute minima at  $x = \pm\sqrt{5/2}$
3. (a)  $A(\theta) = \sin \theta + (\sin \theta)(\cos \theta)$   
(b)  $\theta = \pi/3$   
(c) Since The maximum area is  $\frac{3\sqrt{3}}{4}$
4. Use  $n = 5$ :  $x \approx 1.3247 \dots$
5. (a)  $x_{n+1}x_n - \tan(x_n)$   
(b)  $n = 3$ :  $x \approx 3.14159265$   
(c) nine digits after the decimal point
6. (a)  $n = 5$ :  $x_5 \approx 2.71828183$   
(b) The number  $x_5$  agrees with  $e$  to 6 digits after the decimal point, not just 5 digits.